

Effects of winning cues and relative payout on choice between simulated slot machines

Marcia L. Spetch¹ , Christopher R. Madan² , Yang S. Liu^{1,3}  & Elliot A. Ludvig⁴ 

Department of Psychology, University of Alberta, Edmonton, AB, Canada,¹ School of Psychology, University of Nottingham, Nottingham, UK,² Department of Psychiatry, University of Alberta, Edmonton, AB, Canada³ and Department of Psychology, University of Warwick, Coventry, UK⁴

ABSTRACT

Background and aims Cues associated with winning may encourage gambling. We assessed the effects on risky choice of slot machine of: (1) neutral sounds paired with winning, (2) casino-related cues (such as the sound of coins dropping and pictures of dollar signs) and (3) relative payouts. **Design** Experimental studies in which participants repeatedly chose between safer and riskier simulated slot machines. Safer slot machines paid the same amount regardless of which symbols lined up. Risky machines paid different amounts depending on which symbols lined up. Effects of initially neutral sounds paired with the best payout were assessed between-groups (experiment 1a) and within-participants (experiment 1b). In experiment 2, pairing of casino-related audiovisual cues with payout was assessed within participants, and cue timing was assessed between groups. **Setting** A university research laboratory in Edmonton, Canada. **Participants** Undergraduate students ($n = 630$ across three experiments). **Measurements** Preference for riskier over safer machines, preference between machines that differed in cues, payout recall and frequency estimates for payouts. Risky choice was calculated as the proportion of choices of the risky machine when presented with a fixed machine of the same expected value. **Findings** In experiment 1a, risky choice was slightly increased by pairing a sound with the best payout compared with pairing the sound with a lower payout ($P = 0.04$, $d = 0.28$) but not compared with no sound [$P = 0.36$, $d = 0.13$, Bayes factors $(BF)_{10} = 0.22$]. In experiment 1b, people did not prefer a machine with a best-payout sound over one with a lower-payout sound ($P = 0.67$, $d = 0.03$, $BF_{10} = 0.11$). Relative payout affected choice: risky choices were higher for high- than low-payout decisions ($P < 0.001$, $d = 0.53$). In experiment 2, people preferred machines with casino-related cues paired with winning ($P < 0.001$, $r^2 = 0.11$) and cue timing (at choice or concurrently with the win) had no effect ($P = 0.95$, $r^2 = 0.0$, $BF_{10} = 0.05$). Casino-related cues also enhanced payout memory ($P = 0.013$ and 0.006). Cue effects were not specific to risk: people also preferred fixed-payout machines with casino-related cues ($P < 0.001$, $r^2 = 0.16$). **Conclusions** In a gambling simulation, student participants chose more risky slot machines when payouts were relatively higher and when casino-related cues were associated with payouts. Pairing a neutral sound with the best payout did not consistently affect slot machine choice, and the effect of casino cues did not depend on their timing. Casino-related cues enhanced payout memory.

Keywords Casino cues, gambling, memory, relative payout, risky choice, slot machines, winning sounds.

Correspondence to: Marcia Spetch, Department of Psychology, University of Alberta, P217 Biological Sciences Building, Edmonton, AB, Canada T6G 2E9. E-mail: mspetch@ualberta.ca

Submitted 27 August 2019; initial review completed 18 November 2019; final version accepted 12 February 2020

INTRODUCTION

Cues associated with winning have powerful motivating effects on behavior [1]. Casinos, which are designed to motivate gambling [2], highlight winning through lights and sounds that accompany wins, by announcing jackpots won and by showing photos of lucky winners. The importance of winning cues in promoting gambling is widely

recognized [3,4] and has been investigated in human and animal models of gambling [5–9].

Slot machines are an addictive form of gambling [10], and they provide distinctive and often exciting cues for winning. Current digital machines provide elaborate winning cues, created by teams of professionals including graphic designers [2]. Slot machines also provide these cues at a rapid rate [11]: there are no cards to shuffle, horses to race or other players to bid, so players can

complete a game within seconds. Rapid play has been shown to increase risk-seeking in choice tasks [12]. Slot machines, with elaborate winning cues and high event frequency, have been called the ‘crack cocaine of gambling’ ([10], but see [11]).

Winning cues may enhance the appeal of slot machine gambling via learning mechanisms whereby neutral cues paired with winning become reinforcing and increase gambling [13]. An enhancement of gambling by learned cues is also predicted by incentive sensitization theory [14], wherein Pavlovian associations between cues endow the cues with incentive properties that elicit cravings and motivate reward-seeking. Susceptibility to such ‘cue-triggered wanting’ is predictive of problem gambling [15]. Winning cues may also enhance gambling via memory mechanisms. Specifically, cues present during wins may generate an availability bias [16], whereby large wins are more readily recalled [3,17,18], which could increase the desire to play again.

Few studies have directly tested the hypothesis that cues increase gambling behavior through a learned association with winning. Recently, Cherkasova *et al.* [5] investigated the effect of pairing casino-related audiovisual cues with reward feedback on people’s risk tendencies in the Iowa Gambling Task (IGT), which involves multiple small wins and occasional larger losses, and the Vancouver Gambling Task (VGT), which involves described wins of different probabilities and amounts. The cues did not affect reward-maximizing choices on the IGT, but pairing of audiovisual cues with winning increased risky choice in the VGT. This increase was independent of the expected value of the wins, and both choices and eye movements indicated that the audiovisual cues decreased the influence of reward probability. This study appears to provide the first conclusive experimental evidence that casino-related audiovisual cues paired with winning increases people’s tendency to make riskier choices.

Nevertheless, interesting questions remain. First, does the enhancement depend on use of casino-related cues or would risky choice also be enhanced if the cues were initially neutral, such that reinforcing properties were learned through pairing with wins? In prior work we found that visual cues paired with better wins enhanced risky choice when presented as a priming cue, but we did not test for a general enhancement of risky choice [18]. Secondly, does timing of the cues matter? Would casino-related cues increase risky choice both when they are presented concurrently with winning feedback, and when they are presented predictively, before winning feedback? Thirdly, is the effect of casino-related cues specific to risky choices, or do they enhance attraction to all options?

Our studies expand the empirical investigation of how sensory cues affect risky choice. We investigated whether presenting an initially neutral sound (experiment 1) or a

casino-related audiovisual cue (experiment 2) affected preference between simulated slot machines in a risky-choice task. We manipulated which winning outcomes were accompanied by cues and whether the cue was presented predictively or concurrently with winning. Our study population consisted of undergraduate students drawn from a participant pool that past research indicated should have few people with gambling problems (see Supporting information). We investigated whether cues associated with winning enhance the tendency to make risky choices, potentially increasing the allure of gambling.

Experiment 1 also tested the effect of relative payout value. In risky-choice tasks using images of doors as choice stimuli, we found that people overweight the best and worst outcomes in a decision set, leading to more risk-seeking for decisions that occasionally provide the best outcome [18–21]. People also better recall and sometimes overestimate the frequency of the best and worst outcomes in the task [22,23]. Here, we tested whether people show these same biases with dynamic images of slot machines and whether auditory cues affect these biases.

Our data collection plan, exclusion criteria, experimental methods, research hypotheses and primary analyses were pre-registered on the Open Science Framework (OSF) at

<https://osf.io/s9h8n/>. Materials and data for all experiments are also available on OSF.

GENERAL METHODS

Participants

All research was approved by the University of Alberta Research Ethics Board. Participants were recruited from a psychology participant pool and provided written informed consent. They received course credit and a bonus of up to \$5 (Canadian) depending on points earned.

Materials and design

The risky-choice task presented simulated slot machines on the left and/or right of the screen. The left and right cursor keys were used for choices. Each slot machine had unique images, colors and two distinct reel symbols (Fig. 1). At the start of a trial, the symbols for each machine were offset (Fig. 1: left machine). When a machine was selected, its reels began to spin and then stopped sequentially from left to right, taking 3 s. The spin always ended with three identical symbols lined up, and a 1-s message below the selected machine indicated points won.

The machines differed in average payout (high- or low-value) and constancy of the payout: fixed machines paid the same amount regardless of which symbols lined up, whereas risky machines paid different amounts depending on which symbols lined up; each payout occurred with a




Figure 1 Screen shot showing two slot machines after choice of the right machine was completed. [Colour figure can be viewed at wileyonlinelibrary.com]

50/50 chance. The fixed low-value machine paid 20 points, the risky low-value machine paid 10 or 30 points, the fixed high-value machine paid 60 points and the risky high-value machine paid 50 or 70 points.

Procedure

The choice task had three trial types: single-option trials presented only one machine that had to be selected; these trials ensured exposure to all outcomes, limiting potential avoidance of initially unlucky machines [24]. Catch trials presented a choice between machines that differed in expected payout (high versus low value); these trials provided a manipulation check that participants had learned the contingencies and chose to maximize points/money. As per our standard practice [25,26] and pre-registration, participants who chose reward-maximizing options on fewer than 60% of the catch trials were excluded (see Supporting information). Decision trials presented a choice between fixed and risky machines of equal expected value and provided a measure of risk preference independent of expected-value maximization.

The choice phase was followed by two memory tests. In a recall test, participants were shown each slot machine individually and asked to enter the 'number of points you first think of'; this tested availability biases in memory for payouts. In a frequency-judgment test, each machine was again presented individually with a list of all payouts in the task. Participants were asked to type the percentage of time the machine led to each payout; this assessed distortions in remembered frequency of payouts.

Measures

Risky choice was calculated as the proportion of choices of the risky machine when presented with a fixed machine of

the same expected value. In experiments 1b and 2 we also measured choice between machines that differed in associated cues. In all experiments, we measured payout recall and frequency estimates for each payout.

Statistical analyses

A priori predictions were tested with *t*-tests, with Cohen's *d* providing effect sizes. Corresponding Jeffreys–Zellner–Siow (JZS) Bayes factors (BF_{10}) using a medium prior were calculated [27]. Recall test results were analyzed with χ^2 and McNemar's tests. In experiment 2, Gaussian linear mixed-effects modeling [28], fitted by maximum likelihood, was conducted in R version 3.5.0 [29] to assess both main effects and interactions, with subjects treated as a random effect. For each fixed effect, we report a likelihood-ratio test with 95% confidence interval and effect size as r^2 . Bayes factors showing the relative odds in favor of a model containing the fixed effect against a null model without the effect are also provided.

Additional details are provided in Supporting information and the Methods sections below.

EXPERIMENT 1

This experiment tested the effects of (i) auditory cues associated with winning and (ii) relative payout on choice between risky and safe options. People chose between simulated slot machines in a design similar to our previous studies [22]. Participants experienced the slot machines presented individually and in pairs that provided a risky low-payout, a fixed low-payout, a risky high-payout or a fixed high-payout. In contrast to the IGT, for which choosing the riskiest option provides less reward in the long term, choices here were between machines with the same

expected value but different variability—risk preference was thus measured independently of reward maximization. A preliminary study (see Supporting information) found no evidence that initially neutral sounds presented concurrently with the best and worst payouts affected the tendency to overweight these outcomes.

Experiment 1a tested the effect of an initially neutral auditory cue presented to predictively signal payouts on high-value risky machines. For the group best-cued, the cue occurred before the best payout and, for the group lower-cued, before the lower payout; no cues were presented for the group no-cue. We predicted that the group best-cued would choose the high-value risky option more often than the lower-cued or no-cue groups. Experiment 1b was a within-subject replication in which one high-value risky machine had a predictive sound for the best win and one had a predictive sound for a lower win. We predicted that risky choice would be higher on the best-cued machine than on the lower-cued machine. In both experiments, we also predicted greater risky choice for high- than low-value decisions, consistent with overweighting of the best and worst wins.

METHODS

Participants

In experiment 1a, 328 participants were randomly assigned to three groups. Eighteen were excluded for failing the catch-trial criterion, leaving 310 participants (202 females, 98 males, mean age = 19.3 years). In experiment 1b, 131 participants were tested. Seven failed the catch-trial criterion, leaving 124 participants (79 females, 43 males, mean age = 19.2 years).

Materials and design

Auditory cues were distinct 1.1-s neutral sounds. In experiment 1a, the cue appeared only with the high-value risky machine and preceded the best payout (70) for the group best-cued and the lower payout (50) for the group lower-cued. The group no-cue received no auditory cues. Experiment 1b used a within-subject design with four risky machines. Two visually distinct but functionally equivalent high-value risky machines gave the same payouts, but one had a predictive sound before the best (70) win and the other had a different predictive sound before the lower (50) win. Two visually distinct low-value risky machines were functionally equivalent, yielding 10 or 30 points, and neither had sounds.

RESULTS

In experiment 1a, our prediction that the group best-cued would make more high-value risky choices than the other

two groups was modestly supported for the comparison between the group best-cued and the group lower-cued ($t_{(209)} = 2.07, P = 0.040, d = 0.28, BF_{10} = 1.10$), but not for the comparison between the group best-cued and the group no-cue ($t_{(203)} = 0.91, P = 0.364, d = 0.13, BF_{10} = 0.22$) (see Fig. 2a). As predicted, participants in all groups chose the risky machine significantly more often for high-value than low-value decisions (no-cue, $t_{(98)} = 5.49, P < 0.001, d = 0.55, BF_{10} > 150$; best-cued, $t_{(105)} = 9.45, P < 0.001, d = 0.92, BF_{10} > 150$; lower-cued, $t_{(104)} = 6.23, P < 0.001, d = 0.61, BF_{10} > 150$).

In experiment 1b, contrary to our prediction participants did not prefer the best-cued over the lower-cued machine ($t_{(121)} = 0.33, P = 0.629, d = 0.03, BF_{10} = 0.11$). As predicted, however, participants chose the risky option more for choices between high- than low-value machines (Fig. 2b), whether the high-value risky machine had the best payout cued ($t_{(121)} = 5.89, P < 0.001, d = 0.53,$




Figure 2 Mean risky choice (+95% confidence interval) for experiment 1. (a) In experiment 1a, people selected the risky option more often for high- than low-value decisions, consistent with an overweighting of the extreme payouts, with only small differences between auditory cue groups. (b) In experiment 1b, participants chose the risky option more for high- than for low-value machines, and there was no significant difference between high-value machines that had auditory cues for the best payout or the lower payout. [Colour figure can be viewed at wileyonlinelibrary.com]

$BF_{10} > 150$) or the lower payout cued [$t_{(121)} = 7.61$, $P < 0.001$, $d = 0.69$, $BF_{10} > 150$].

On recall tests of experiment 1a (Fig. 3a), more people in all groups reported the best payout (70) than the lower payout (50) for the high-value risky machine (no-cue $\chi^2_{(1, n = 88)} = 22.0$, $P < 0.001$, $BF_{10} > 150$; best-cued, $\chi^2_{(1, n = 97)} = 8.67$, $P = 0.003$, $BF_{10} = 15.1$; lower-cued, $\chi^2_{(1, n = 95)} = 8.85$, $P = 0.003$, $BF_{10} = 16.5$) and more people reported the worst payout (10) than the higher payout (30) for the low-value risky machine (no-cue, $\chi^2_{(1, n = 87)} = 12.5$, $P < 0.001$, $BF_{10} = 100.2$; best-cued, $\chi^2_{(1, n = 96)} = 24.0$, $P < 0.001$, $BF_{10} > 150$; lower-cued, $\chi^2_{(1, n = 94)} = 26.6$, $P < 0.001$, $BF_{10} > 150$). The groups did not differ in their recall for either the low-value ($\chi^2_{(2, n = 277)} = 1.49$, $P = 0.47$, $BF_{10} = 0.05$) or high-value risky machine ($\chi^2_{(2, n = 280)} = 2.73$, $P = 0.26$, $BF_{10} = 0.1$). In experiment 1b (Fig. 3b), more people reported 70 than 50 for both high-value machines (best-cued: $\chi^2_{(1)} = 15.09$, $P < 0.001$, $BF_{10} > 150$; lower-cued: $\chi^2_{(1)} = 6.81$, $P = 0.009$, $BF_{10} = 5.98$).

Reporting differences were small ($\chi^2_{(1)} = 4.05$, $P = 0.044$, $BF_{10} = 1.58$) or not significant ($\chi^2_{(1)} = 2.23$, $P = 0.136$, $BF_{10} = 0.69$) for the low-value risky machines.

Frequency estimates were similar across groups and experiments and showed overweighting of the worst but not the best payouts. Consistent with previous research [22,23], risky choice correlated with memory: stronger memories for the highest payout correlated with more risk-seeking for high-value choices and stronger memories for the lowest payout correlated with risk aversion on low-value choices. These results are reported in Supporting information.

DISCUSSION

In both experiments, auditory cues had no systematic effects on risky choice, but people overweighted the best and worst payouts. This overweighting extends the previously reported extreme-outcome effect [25] to choices between simulated slot machines. Although experiment 1a




Figure 3 Recall test results showing the proportion of participants who reported the worst payout for low-value risky machines and the best payout for high-value risky machines. NC = no-cue, BC = best-cued, LC = lower-cued. (a) Results for the three groups in experiment 1a. (b) Results for each risky machine in experiment 1b. Both low-value risky machines are labelled NC because neither had cues. [Colour figure can be viewed at wileyonlinelibrary.com]

showed a mild difference between cuing the best and lower wins, the effect was smaller than the relative payout effect and did not replicate in experiment 1b. Thus, cues with no prior winning connotation did not reliably enhance risky choice when paired predictively with the best win.

EXPERIMENT 2

Experiment 2 tested whether an audiovisual cue that already had winning connotations (casino sounds and money-related visual stimuli), presented either concurrently or predictively with winning, would increase risky choice, consistent with Cherkasova *et al.* [5]. We tested the specificity of cuing effects by comparing the effect of cues on both risky and fixed machines.

METHODS

Participants

We randomly assigned 258 participants to group concurrent and group predictive. Sixty-two participants were excluded for failing the catch-trial criterion, leaving 196 participants (152 females, 42 males, mean age = 19.2 years).

Materials, design and procedure

Winning cues were casino sounds (such as coins dropping) accompanied by dollar signs or gold bars displayed on the chosen machine. All stimuli are available on the OSF repository. For the group predictive, the cue was presented when the machine was selected and lasted for 2.6 s while the reels were spinning. For the group concurrent, the cue was presented when the reels stopped spinning and occurred for 2.6 s while the winning message was displayed.

There were two fixed machines that always paid 30 points: one had casino-related cues (fixed-CC) and one had no cues (fixed-NC). There were four risky machines. Two risky machines both paid 10 or 50 points with equal probability; one provided casino-related cues (risky-CC) for the 50-point win and the other machine provided no cues (risky-NC). The other risky machines also paid 10 or 50 points but had different expected payouts and were included to motivate selective choice. Machine risky-20 paid 50 with a 20% probability and 10 otherwise, and machine risky-80 paid 50 with an 80% probability and 10 otherwise; neither machine had audiovisual cues. Memory recall and frequency judgments were tested for all machines.

RESULTS

Figure 4a shows that participants in both groups chose the risky-CC machine more often than the risky-NC machine.




Figure 4 Mean (+95% confidence interval) proportion of trials on which machines with or without casino cues were chosen by the two groups in experiment 2. (a) Risky machines. (b) Fixed machines. [Colour figure can be viewed at wileyonlinelibrary.com]

A likelihood ratio test, with a between-group factor of timing (concurrent or predictive) and a within-subject factor of cue (risky-CC or risky-NC), confirmed a significant main effect of cue ($\chi^2_{(1)} = 22.53$, $P < 0.001$, $r^2 = 0.11$, $BF_{10} > 150$), but no effect of timing ($\chi^2_{(1)} = 0.004$, $P = 0.95$, $r^2 = 0.0$, $BF_{10} = 0.05$), and no interaction ($\chi^2_{(1)} = 2.43$, $P = 0.12$, $r^2 = 0.01$, $BF_{10} = 0.17$).

Participants also chose the fixed-CC machine more often than the fixed-NC machine (Fig. 4b). There was a significant main effect of cue ($\chi^2_{(1)} = 34.51$, $P < 0.001$, $r^2 = 0.16$, $BF_{10} > 150$) and a small interaction between cue and timing ($\chi^2_{(1)} = 4.22$, $P = 0.04$, $r^2 = 0.02$, $BF_{10} = 0.42$), due to a slightly larger effect of the cue in the predictive group, but there was no main effect of timing ($\chi^2_{(1)} = 0.16$, $P = 0.69$, $r^2 = 0.0$, $BF_{10} = 0.05$).

On recall tests (Fig. 5a), more people in both groups reported the higher payout (50) for the risky-CC machine than the risky-NC machine. McNemar's test for related samples, using only participants who reported either 10 or 50 for both risky machines, showed significant effects for both groups (concurrent, $\chi^2_{(1)} = 6.15$, $P = 0.013$; predictive, $\chi^2_{(1)} = 7.41$, $P = 0.006$).




Figure 5 Recall test results for experiment 2 showing proportion of participants who reported the best payout on the risky machine with no cue (NC) or with a casino cue (CC). [Colour figure can be viewed at wileyonlinelibrary.com]

As presented in Supporting information, frequency judgments for the best win were also higher for the risky-CC machine than the risky-NC machine for both groups. Moreover, memory for the better win correlated positively with risky choice for both groups.

GENERAL DISCUSSION

We showed that preference in a risky-choice task was increased by casino-related cues, but not by neutral cues paired with winning in the task. Specifically, in experiment 1, with more than 400 participants, we found strong and consistent effects of relative payout but no or weak effects of pairing neutral sounds with winning on preference. In experiment 2, casino-related cues, presented predictively or concurrently, increased machine preference. This effect, however, was not specific to risky choices or cue timing. People preferred both risky and fixed machines that had casino-related cues, regardless of cue timing.

The robust effect of casino cues on slot machine preference in experiment 2 is consistent with the findings of Cherkasova *et al.* [5]. As in their study, people preferentially chose a risky machine if wins were accompanied by casino sounds and win-related visual cues. Our study extends their findings in three ways. First, the effect of casino cues was not specific to choices involving risk because the cues also enhanced preference for machines that provided the same win every time. Casino-related cues thus appear to have a general attraction effect. Secondly, casino-related cues had the same effect whether they were presented predictively, during the spinning of the reels, or concurrently with the win. Thus, cue timing does not seem to matter. Thirdly, casino-related cues affected memory for wins. Cues associated with the best win increased recall of the best win and estimates of how often the best win occurred.

Like Cherkasova *et al.* [5], we tested a population that was unlikely to include many problem gamblers. One limitation, therefore, is that our results may apply best to young people with little slot machine experience. Indeed, our findings may be most relevant to understanding the allure of slot-machine gambling during initial exposure, whether in physical casinos or on-line gaming sites. If casino-related cues enhance attraction to machines and makes wins more memorable, this could encourage continued play, thereby providing more exposure and opportunity for gambling addictions to develop. There is evidence suggesting that exposure to free play can increase subsequent gambling with real money [30] and that exposure to simulated internet gambling may provide a gateway for transition to real-money gambling [31,32].

Similar to many laboratory studies that have provided insights into gambling behavior [6,33,34], another limitation is that our task differed from real-world slot machines in payout structure and lack of requirement to bet with money. Our results may therefore be most relevant to understanding what makes certain machines more attractive than others. Whether susceptibility to the allure of casino-related cues predicts gambling persistence or the development of gambling addition is an important direction for future research.

The observed relative payout effect is consistent with recent evidence that people overweight outcomes near the edges of an experienced distribution, increasing avoidance of options that could lead to the worst outcome and enhancing attraction to options that could lead to the best outcome ([19,22,35]; see [26] for a review). People also showed enhanced recall for the best and worst payouts. Associating a neutral sound with the best payout, however, did not substantially affect payout memory or risky choice.

Despite limitations, our results have implications for real-world gambling. First, they suggest that people are attracted to machines that provide casino-related cues, but this attraction is not specific to when the cues occur, nor how risky the machine is. Secondly, pairing neutral sounds with winning may not enhance gambling preferences, at least with short exposures. Thirdly, the relative payout effect suggests that the best and worst payouts of a gambling experience may disproportionately affect behavior. In studies using doors as choice stimuli, we found that relative payout effects were specific to a particular episode and set of cues [36]. Therefore, in a casino, overweighting of the best and worst payout is probably based on a specific gambling episode rather than the entire set of gambling experiences.

Finally, we found that relative payout and casino-related cues affect not only choice behavior but also memory for the wins. This result is consistent with findings that casino sounds increase peoples' estimates of how often they won [8]. An important future research direction is to determine whether relative payout and casino-related cues affect not only risk preference in the moment, but also attraction to slot machines in the future.

Declaration of interests

M.L.S. was the Principal Investigator (PI) and E.L. and C.M. were co-PIs on research grants funded by the Alberta Gambling Research Institute that ended in February 2019.

Acknowledgements

This research was supported by grants from the Natural Sciences and Engineering Research Council of Canada (04133) and the Alberta Gambling Research Institute (74) held by M.L.S. We thank Jeffrey Pisklak for assistance with statistical analyses and Cody Freas, Juliana Montoya, Maria Shanks and Josh Yong for assistance in testing participants.

References

1. Robinson T. E., Yager L. M., Cogan E. S., Saunders B. T. On the motivational properties of reward cues: individual differences. *Neuropharmacology* 2014; **76**: 450–9.
2. Schull N. D. *Addiction by Design: Machine Gambling in Las Vegas*. Princeton, NJ: Princeton University Press; 2012.
3. Griffiths M. Fruit machine gambling: the importance of structural characteristics. *J Gambl Stud* 1993; **9**: 101–20.
4. Wulfert E., Maxson J., Jardin B. Cue-specific reactivity in experienced gamblers. *Psychol Addict Behav* 2009; **23**: 731–5.
5. Cherkasova M. V., Clark L., Barton J. J., Schulzer M., Shafiee M., Kingstone A. *et al.* Win-concurrent sensory cues can promote riskier choice. *J Neurosci* 2018; **38**: 10362–70.
6. Barrus M. M., Winstanley C. A. Dopamine D3 receptors modulate the ability of win-paired cues to increase risky choice in a rat gambling task. *J Neurosci* 2016; **36**: 785–94.
7. Crockford D. N., Goodyear B., Edwards J., Quickfall J., el-Guebaly N. Cue-induced brain activity in pathological gamblers. *Biol Psychiatry* 2005; **58**: 787–95.
8. Dixon M. J., Harrigan K. A., Santesso D. L., Graydon C., Fugelsang J. A., Collins K. The impact of sound in modern multiline video slot machine play. *J Gambl Stud* 2014; **30**: 913–29.
9. Marshall A. T., Kirkpatrick K. Reinforcement learning models of risky choice and the promotion of risk-taking by losses disguised as wins in rats. *J Exp Psychol* 2017; **43**: 262–79.
10. Breen R. B., Zimmerman M. Rapid onset of pathological gambling in machine gamblers. *J Gambl Stud* 2002; **18**: 31–43.
11. Dowling N., Smith D., Thomas T. Electronic gaming machines: are they the 'crack-cocaine' of gambling? *Addiction* 2005; **100**: 33–45.
12. Madan C. R., Spetch M. L., Ludvig E. A. Rapid makes risky: time pressure increases risk seeking in decisions from experience. *J Cogn Psychol* 2015; **27**: 921–8.
13. Skinner B. F. *Science and Human Behavior*. New York: The Free Press; 1953.
14. Robinson T. E., Berridge K. C. The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res Rev* 1993; **18**: 247–91.
15. Davey B., Cummins R. Testing an incentive-sensitisation approach to understanding problem slot-machine gambling using an online slot-machine simulation. *J Gambl Stud* 2018; **34**: 773–84.
16. Tversky A., Kahneman D. Availability: a heuristic for judging frequency and probability. *Cogn Psychol* 1973; **5**: 207–32.
17. Griffiths M. D. The role of cognitive bias and skill in fruit machine gambling. *Br J Psychol* 1994; **85**: 351–69.
18. Ludvig E. A., Madan C. R., Spetch M. L. Priming memories of past wins induces risk seeking. *J Exp Psychol Gen* 2015; **144**: 24–9.
19. Ludvig E. A., Madan C. R., Spetch M. L. Extreme outcomes sway risky decisions from experience. *J Behav Decis Mak* 2014; **27**: 146–56.
20. Ludvig E. A., Madan C. R., Pisklak J. M., Spetch M. L. Reward context determines risky choice in pigeons and humans. *Biol Lett* 2014; **10**: 20140451.
21. Ludvig E. A., Madan C. R., McMillan N., Xu Y., Spetch M. L. Living near the edge: how extreme outcomes and their neighbors drive risky choice. *J Exp Psychol Gen* 2018; **147**: 1905–18.
22. Madan C. R., Ludvig E. A., Spetch M. L. Remembering the best and worst of times: memories for extreme outcomes bias risky decisions. *Psychon Bull Rev* 2014; **21**: 629–36.
23. Madan C. R., Ludvig E. A., Spetch M. L. The role of memory in distinguishing risky decisions from experience and description. *Q J Exp Psychol* 2017; **70**: 2048–59.
24. Denrell J., March J. G. Adaptation as information restriction: the hot stove effect. *Org Sci* 2001; **12**: 523–38.
25. Ludvig E. A., Spetch M. L. Of black swans and tossed coins: is the description-experience gap in risky choice limited to rare events? *PLOS ONE* 2011; **6**: e20262.
26. Madan C. R., Ludvig E. A., Spetch M. L. Comparative inspiration: from puzzles with pigeons to novel discoveries with humans in risky choice. *Behav Processes* 2019; **160**: 10–10.
27. Morey R. D., Rouder J. N. Bayes factor approaches for testing interval null hypotheses. *Psychol Methods* 2011; **16**: 406–19.
28. Pinheiro J., Bates D., DebRoy S., Sarkar D., R Core Team. nlme: Linear and nonlinear mixed effects models. 2018. Available

- at: <https://CRAN.R-project.org/package=nlme> (R package version 3.1-137) (accessed 6 August 2019).
29. R Core Team. R: A language and environment for statistical computing. Vienna, Austria. 2018. Available at: <https://www.R-project.org/> (accessed 6 August 2019)
 30. Bednarz J., Delfabbro P., King D. Practice makes poorer: practice gambling modes and their effects on real-play in simulated roulette. *Int J Ment Health Add* 2013; **11**: 381–95.
 31. Hayer T., Kalke J., Meyer G., Brosowski T. Do simulated gambling activities predict gambling with real money during adolescence? Empirical findings from a longitudinal study. *J Gambl Stud* 2018; **34**: 929–47.
 32. Armstrong T., Rockloff M., Browne M., Li E. An exploration of how simulated gambling games may promote gambling with money. *J Gambl Stud* 2018; **34**: 1165–84.
 33. Clark L., Crooks B., Clarke R., Aitken M. R., Dunn B. D. Physiological responses to near-miss outcomes and personal control during simulated gambling. *J Gambl Stud* 2012; **28**: 123–37.
 34. Olsen C. M., Winder D. G. Operant sensation seeking engages similar neural substrates to operant drug seeking in C57 mice. *Neuropsychopharmacology* 2009; **34**: 1685–94.
 35. Konstantinidis E., Taylor R. T., Newell B. R. Magnitude and incentives: revisiting the overweighting of extreme events in risky decisions from experience. *Psychon Bull Rev* 2018; **25**: 1925–33.
 36. Madan C. R., Spetch M. L., Machado F. M. S., Mason A. M. A., Ludvig E. A. Encoding context determines risky choice. *PsyArXiv* 2019. Available at: <https://doi.org/10.31234/osf.io/cujqf> (accessed 31 January 2020).

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Table S1 Payouts and cue conditions for the six slot machines used in Experiment 2.

Figure S1 Risky choice (mean and 95% CI) for the Preliminary Experiment. A. Participants in all three groups selected the risky option more often in the high-value decisions than in the low-value decisions, consistent with an overweighting of the extreme outcomes, but there were only small differences between the groups based on the cues.

Figure S2 Memory test results for the Preliminary Experiment. A. Proportion of participants (based on those who reported a valid payout) who reported the worst payout for the low-value risky machine and the best payout for the high-value risky machine. Groups: NC=No Cue, EX = Extreme, N-EX = Non-Extreme. B. Mean reported frequency of the two payouts on each of the risky machines for the three groups. Error bars are 95% confidence intervals.

Figure S3 Mean accuracy on catch trials and risky choice on high-value and low-value decisions as a function of trial block for the three groups in Experiment 1a. Error bars are 95% confidence intervals.

Figure S4 Mean reported frequency of the two payouts on low-value and high-value risky machines for the three groups in Experiment 1a. Error bars are 95% confidence intervals.

Figure S5 Mean reported frequency of the two payouts on each of the two low-value risky machines (neither had a cue) and on the best-cued and lower-cued high-value risky machines in Experiment 1B. Error bars are 95% confidence intervals.

Figure S6 Mean reported frequency of the best payout for the Risky-NC and Risky-CC machines for the two groups in Experiment 2. NC = no cue. CC = casino cue. Error bars are 95% confidence intervals.